

Title	On Laplace transforms of certain special functions (Study on Non-Analytic and Univalent Functions and Applications)
Author(s)	Takano, Katsuo
Citation	数理解析研究所講究録 (2009), 1626: 122-126
Issue Date	2009-01
URL	http://hdl.handle.net/2433/140309
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

On Laplace transforms of certain special functions

高野勝男 [Takano Katsuo]
茨城大学 [Ibaraki University]

1 Introduction

In this note we consider a Laplace transform of the following multiple integral

$$\begin{aligned} g(d; n; y) &= c\pi^{d/2} \frac{1}{y^{\{n(d+1)-d\}/2+1}} \exp\left\{-\frac{a_n^2}{y}\right\} \\ &= \int \cdots \int_{0 \leq v_1 + \cdots + v_{n-1} \leq 1; 0 \leq v_j} \exp\left\{\sum_{j=1}^{n-1} (a_n^2 - a_j^2) v_j / y\right\} \\ &\quad \left[v_1 \cdots v_{n-1} (1 - \sum_{j=1}^{n-1} v_j)\right]^{(d+1)/2-1} dv_1 \cdots dv_{n-1} \end{aligned} \quad (1)$$

where c is a constant. This multiple integral is coming from the normed product of multidimensional Cauchy densities, namely

$$f(d; a_1, \dots, a_n; x) = \frac{c_1}{\pi_{j=1}^n (a_j^2 + |x|^2)^{(d+1)/2}},$$

where c_1 is the normalised constant and $0 < a_j$; $a_j \neq a_k$ if $j \neq k$, and $x = (x_1, \dots, x_d) \in \mathbf{R}^d$. We note that the density function $g(d; 1; u)$ can be expressed by a confluent hypergeometric function,

$$\begin{aligned} g(d; 1; u) &= \frac{c\pi^{d/2} \Gamma((d+1)/2)^2}{\Gamma(d+1)} \\ &\quad \cdot \frac{1}{u^{d/2+2}} \exp\left\{-\frac{b^2}{u}\right\} {}_1F_1\left(\frac{d+1}{2}; d+1; \frac{(b^2 - a^2)}{u}\right) \end{aligned} \quad (2)$$

for all dimension d . For the dimension $d = 3$ we obtain

$$\begin{aligned} g(3; 1; u) &= \frac{c\pi^{3/2}}{(b^2 - a^2)^2} \left[\frac{1}{u^{3/2}} (\exp\{-\frac{a^2}{u}\} + \exp\{-\frac{b^2}{u}\}) \right. \\ &\quad \left. - \frac{2}{(b^2 - a^2)u^{1/2}} (\exp\{-\frac{a^2}{u}\} - \exp\{-\frac{b^2}{u}\}) \right] \end{aligned} \quad (3)$$

and for the dimension $d = 5$ we obtain

$$\begin{aligned}
 g(5; 1; u) = & \frac{c\pi^{5/2}}{(b^2 - a^2)^3} \left[\frac{2!}{u^{3/2}} (\exp\{-\frac{a^2}{u}\} - \exp\{-\frac{b^2}{u}\}) \right. \\
 & - \frac{2 \cdot 3!}{(b^2 - a^2)u^{1/2}} (\exp\{-\frac{a^2}{u}\} + \exp\{-\frac{b^2}{u}\}) \\
 & \left. + \frac{4!u^{1/2}}{(b^2 - a^2)^2} (\exp\{-\frac{a^2}{u}\} - \exp\{-\frac{b^2}{u}\}) \right]. \quad (4)
 \end{aligned}$$

2 The Laplace transforms of the case $n = 1$

Let us denote the Laplace transform of the functions $g(d; 1; u)$ by $\eta(d; 1; s)$ and let us take a branch of the value $\eta(d; +0) = 1$. We obtain

$$\begin{aligned}
 \eta(3; 1; s) = & \frac{c\pi^2}{(b^2 - a^2)^2} \left\{ \left(\frac{1}{a} \exp\{-2a\sqrt{s}\} + \frac{1}{b} \exp\{-2b\sqrt{s}\} \right) \right. \\
 & \left. - \frac{2}{(b^2 - a^2)\sqrt{s}} (\exp\{-2a\sqrt{s}\} - \exp\{-2b\sqrt{s}\}) \right\}, \quad (5)
 \end{aligned}$$

for the dimension $d = 3$ and for the dimension $d = 5$ we obtain

$$\begin{aligned}
 \eta(5; 1; s) = & \frac{c\pi^3}{(b^2 - a^2)^3} \left[2! \left(\frac{1}{a} \exp\{-2a\sqrt{s}\} - \frac{1}{b} \exp\{-2b\sqrt{s}\} \right) \right. \\
 & - \frac{2 \cdot 3!}{(b^2 - a^2)\sqrt{s}} (\exp\{-2a\sqrt{s}\} + \exp\{-2b\sqrt{s}\}) \\
 & + \frac{4!}{(b^2 - a^2)^2 s} \left\{ a \exp\{-2a\sqrt{s}\} \left(1 + \frac{1}{2a\sqrt{s}} \right) \right. \\
 & \left. \left. - b \exp\{-2b\sqrt{s}\} \left(1 + \frac{1}{2b\sqrt{s}} \right) \right\} \right]. \quad (6)
 \end{aligned}$$

Making use of the formula

$$K_\nu(z) = \frac{1}{2} \left(\frac{1}{2} z \right)^\nu \int_0^\infty \exp\left\{-t - \frac{z^2}{4t}\right\} \frac{dt}{t^{\nu+1}}, \quad (7)$$

we obtain Laplace transforms for the odd dimensional case,

$$\begin{aligned}
 \eta(d; 1; s) = & \frac{c\pi^{d/2}}{\{(b^2 - a^2)^{l+1}\} \sum_{j=0}^l \frac{(-1)^{2l-j}(l+j)!}{(b^2 - a^2)^j} \binom{l}{j} 2^{j+1/2}} \cdot \\
 & \left\{ a^{2j-1} \frac{K_{(j-1)+1/2}(2a\sqrt{s})}{(2a\sqrt{s})^{j-1+1/2}} + (-1)^{l+1+j} b^{2j-1} \frac{K_{(j-1)+1/2}(2b\sqrt{s})}{(2b\sqrt{s})^{j-1+1/2}} \right\}, \quad (8)
 \end{aligned}$$

where

$$K_{n+1/2}(z) = \left(\frac{\pi}{2z} \right)^{1/2} \exp\{-z\} \sum_{r=0}^n \frac{(n+r)!}{r!(n-r)!(2z)^r} \quad (9)$$

(cf. G. Watson [7]).

3 Nonzero region of the Laplace transform

In general, we obtain

$$\begin{aligned} \eta(d; 1; s) &= c\pi^{l+1} \int_0^1 \frac{v^l(1-v)^l}{(b^2 - (b^2 - a^2)v)^{(d+2)/2}} \\ &\exp\{-2(b^2 - (b^2 - a^2)v)^{1/2}\sqrt{s}\} \\ &\sum_{r=0}^{l+1} \frac{(l+1+r)!}{r!(l+1-r)!2^r} \{(b^2 - (b^2 - a^2)v)^{1/2}\sqrt{s}\}^{l+1-r} dv \end{aligned} \quad (10)$$

for the odd dimension $d = 2l + 1$ and we see that $\eta(d; 1; s) \neq 0$ in a neighborhood of the origin except at the origin and $\eta(d; 1; s)$ is bounded in a neighborhood of the origin. From the above Laplace transforms (8) we can see that the Laplace transforms $\eta(d; 1; s)$ do not vanish in a neighborhood of the infinity point.

4 The Laplace transform for the general case

Let us obtain the Laplace transform of $g(d; n; y)$ for the odd dimension. The following equality holds:

$$\begin{aligned} &\exp\{-a_n^2 t\} \cdot t^{nl+n-1} \\ &\cdot \int \cdots \int_{0 \leq v_1 + \cdots + v_{n-1} \leq 1; 0 \leq v_j} \exp\left\{\sum_{j=1}^{n-1} (a_n^2 - a_j^2)tv_j\right\} \\ &\cdot [v_1 \cdots v_{n-1} (1 - \sum_{j=1}^{n-1} v_j)]^l dv_1 \cdots dv_{n-1} \\ &= (-1)^{(l+1)(n-1)} \sum_{r=1}^n \exp\{-a_r^2 t\} \\ &\cdot \left[\sum_{j_{n-1}=0}^l \frac{l!(l+j_{n-1})!}{j_{n-1}!(l-j_{n-1}+1)!} \frac{1}{B(r; n, n-1)^{l+j_{n-1}+1}} \right. \\ &\cdot \sum_{j_{n-2}=0}^{l-j_{n-1}} \frac{(l-j_{n-1})!(l+j_{n-2})!}{j_{n-2}!(l-j_{n-1}-j_{n-2})!} \frac{1}{B(r; n, n-2)^{l+j_{n-2}+1}} \\ &\cdot \sum_{j_{n-3}=0}^{l-j_{n-1}-j_{n-2}} \frac{(l-j_{n-1}-j_{n-2})!(l+j_{n-3})!}{j_{n-3}!(l-j_{n-1}-j_{n-2}-j_{n-3})!} \\ &\cdot \frac{1}{B(r; n, n-3)^{l+j_{n-3}+1}} \cdots \\ &\cdot \sum_{j_1=0}^{l-j_{n-1}-j_{n-2}-\cdots-j_2} \frac{(l-j_{n-1}-j_{n-2}-\cdots-j_2)!(l+j_1)!}{j_1!(l-j_{n-1}-j_{n-2}-\cdots-j_2-j_1)!} \\ &\cdot \left. \frac{1}{B(r; n, 1)^{l+j_1+1}} t^{l-j_1-\cdots-j_{n-2}-j_{n-1}} \right] \end{aligned} \quad (11)$$

where $B(r; n, 1) = a_r^2 - a_{r+1}^2, \dots, B(r; n, n-r) = a_r^2 - a_n^2, B(r; n, n-r+1) = a_r^2 - a_1^2, \dots, B(r; n, n-1) = a_r^2 - a_{r-1}^2$.

Theorem 1 *The Laplace transform is given in the following form.*

$$\begin{aligned}
 \eta(d; n; s) &= \int_0^\infty \exp\{-sy\} g(d; n; y) dy \\
 &= c_1 \pi^{l+1/2} (-1)^{(l+1)(n-1)} \\
 &\quad \sum_{r=1}^n \left[\sum_{j_{n-1}=0}^l \frac{l!(l+j_{n-1})!}{j_{n-1}!(l-j_{n-1}+1)!} \frac{1}{B(r; n, n-1)^{l+j_{n-1}+1}} \right. \\
 &\quad \sum_{j_{n-2}=0}^{l-j_{n-1}} \frac{(l-j_{n-1})!(l+j_{n-2})!}{j_{n-2}!(l-j_{n-1}-j_{n-2})!} \frac{1}{B(r; n, n-2)^{l+j_{n-2}+1}} \\
 &\quad \sum_{j_{n-3}=0}^{l-j_{n-1}-j_{n-2}} \frac{(l-j_{n-1}-j_{n-2})!(l+j_{n-3})!}{j_{n-3}!(l-j_{n-1}-j_{n-2}-j_{n-3})!} \\
 &\quad \frac{1}{B(r; n, n-3)^{l+j_{n-3}+1}} \cdots \\
 &\quad \sum_{j_1=0}^{l-j_{n-1}-j_{n-2}-\cdots-j_2} \frac{(l-j_{n-1}-j_{n-2}-\cdots-j_2)!(l+j_1)!}{j_1!(l-j_{n-1}-j_{n-2}-\cdots-j_2-j_1)!} \\
 &\quad \left. \frac{1}{B(r; n, 1)^{l+j_1+1}} 2 \left(\frac{2}{2a_r \sqrt{s}} \right)^{\sum_{p=1}^{n-1} j_p - 1/2} K_{\sum_{p=1}^{n-1} j_p - 1/2} (2a_r \sqrt{s}) \right] \quad (12)
 \end{aligned}$$

Here $K_{\sum_{p=1}^{n-1} j_p - 1/2} (2a_r \sqrt{s})$ is given by (9).

References

- [1] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions, New York, Dover, 1970.
- [2] D. H. Kelker, Infinite divisibility and variance mixtures of the normal distribution, *Ann. Math. Statist.*, 42 (1971), 802–808.
- [3] G. Sansone & J. Gerretsen, Lectures on the theory of functions of a complex variable, 1. Holomorphic functions, P. Noordhoff-Groningen, 1960
- [4] K. Takano, On Laplace transforms of some probability densities, *Applied Mathematics and computations*, 2004.
- [5] ———, Hypergeometric functions and infinite divisibility of probability distributions consisting of Gamma functions, *International J. Pure and Applied Math.*, 20 no.3 (2005), 379-404.
- [6] ———, On Laplace transforms of certain confluent hypergeometric functions, Proceedings of international symposium on new development of geometric function theory and its applications, Edited by Maslina Darus and Sigeyoshi Owa, School of Mathematical Sciences, Faculty of Science & Technology, Universiti Kebangsaan Malaysia, 2008, 260-264.
- [7] G. N. Watson, a treatise on the THEORY OF BESSEL FUNCTIONS, Cambridge University Press, Second edition, Reprinted 1980.

Katsuo TAKANO
 Mathematics, Faculty of Science
 Ibaraki University
 E-mail: ktaka@mx.ibaraki.ac.jp